

## Restricted Medication List on Formweb

| Restricted Medications      |                                  | Restrictions  |
|-----------------------------|----------------------------------|---|
| Antipsychotic               | Aripiprazole (Abilify)           | Restricted to psychiatry or continuation of home meds   |
|                             | Asenapine (Saphris)              | Restricted to psychiatry or continuation of home meds   |
|                             | Clozapine (Clozaril)             | Restricted to psychiatry or continuation of home meds   |
|                             | Lurasidone (Latuda)              | Restricted to psychiatry or continuation of home meds   |
|                             | Paliperidone (Invega)            | Restricted to psychiatry or continuation of home meds   |
| Antibiotics and Antifungals | Ceftaroline fosamil (Teflaro)    | Restricted to use by Infectious Diseases physicians only  |
|                             | Ceftazidime                      | Restricted to use by Infectious Diseases or Nephrology providers only   |
|                             | Ceftazidime/avibactam (Avycaz)   | Restricted to Infections Disease physicians for the treatment of carbapenemase-resistant Enterobacteriaceae and the treatment of multi-drug resistant Pseudomonas resistant to ceftolozane/tazobactam |
|                             | Ceftolozane/tazobactam (Zerbaxa) | Restricted to Infectious Disease physicians for treatment of multi-drug resistant Pseudomonas aeruginosa infections   |
|                             | Cefotaxime (Claforan)            | Restricted to pediatric patients  |

|                            |                            |  |
|----------------------------|----------------------------|--|
|                            | Eravacycline (Xerava)      | Restricted to Infectious Diseases providers or Critical Care providers for use in the following conditions: <ul style="list-style-type: none"> <li>• ESBL/MDR coverage and contraindication to carbapenems</li> <li>• Penicillin-allergic patients with suspected/confirmed MDR infections</li> <li>• MDR <i>Acinetobacter</i> (resistant to cefepime, ampicillin-sulbactam, and meropenem)</li> </ul>   |
|                            | Fidaxomicin (Dificid)      | Restricted to use of recurrent CDI or vancomycin allergy   |
|                            | Fosfomycin (Monurol)       | Restricted to use in treating UTI's with VRE or ESBL-producing organisms   |
|                            | Micafungin (Mycamine)      | Restricted to use by Infectious Diseases or Critical Care physicians   |
|                            | Posaconazole (Noxafil)     | Restricted to adult patients and must be ordered by ID or oncology physicians  |
|                            | Tigecycline (Tygacil)      | Restricted for use in patients with anaphylactic allergy or resistance to all other appropriate available antibiotic options   |
| Analgesics and Anesthetics | Acetaminophen IV (Ofirmev) | Restrictions are: <ul style="list-style-type: none"> <li>• 1x dose in PACU to treat pain in patients meeting criteria per the OSA risk assessment screening tool</li> <li>• Control of fever in neutropenic patients who are unable to receive oral/rectal acetaminophen or IV ketorolac</li> <li>• Abdominal surgery patients at least 70 years or BMI at least 40</li> <li>• Robotic Heart surgery patients: one dose post-op only if the first dose given in the intraoperative period. Must be ordered by a CVT surgeon and ordered through order set</li> <li>• Thoracic surgery patients: 6 doses post-op only if the first dose is given in the intraoperative period. Dose must be ordered by a CVT surgeon.</li> <li>• Colorectal surgery: Acetaminophen injection to allow 48-hour dosing strategy at q6h intervals intraoperatively for a total of 8 doses</li> </ul> |

|                           |                                 |  |
|---------------------------|---------------------------------|--|
|                           | Bupivacaine liposomal (Exparel) | Restricted to adults undergoing anorectal surgeries, colectomies, knee/hip replacement, fractures, and TAVR  |
|                           | Meperidine (Demerol)            | Restricted to the prevention and treatment of drug-induced or blood product-induced rigors and treatment of postoperative shivering  |
|                           | Ketamine oral                   | Restrictions for oral ketamine: <ul style="list-style-type: none"> <li>• Must be prescribed by a palliative care physician and patient must be DNR</li> <li>• Patients must be on 5W, RCU, SPCU and at least 18 years old</li> </ul> |
| Monoclonal antibody       | Adalimumab (Humira)             | Restricted to the use only as induction therapy for adult and peds with ulcerative colitis or crohn's disease  |
|                           | Infliximab (Remicade)           | Restricted to outpatient except for: <ul style="list-style-type: none"> <li>• Inpatient use by gastroenterologists in patients with Crohn's disease or ulcerative colitis who are nearing obstruction</li> </ul>                     |
|                           | Caplacizumab (Cablivi)          | Restricted to hematology/oncology and nephrology providers only  |
| Gastrointestinal Agent    | Methylnaltrexone (Relistor)     | Restricted to use in palliative care and oncology patients that are NPO  |
|                           | Naloxegol (Movantik)            | Restricted to use in patients with opioid induced constipation that have failed 2 or more laxatives  |
| Colony Stimulating Factor | Filgrastim (Neupogen)           | Restricted to pediatric patients requiring doses less than 180mcg  |
| Antiemetic                | Aprepitant (Emend)              | Oral: restricted to use by medical oncologists for the prevention of acute and delayed emesis for moderately or highly emetogenic chemotherapy   |

|                                   |   |   |
|-----------------------------------|---|---|
| Antimigraine                      | Rizatriptan (Maxalt)                      | Restricted to patients intolerant to sumatriptan  |
| Reversal Agents                   | Sugammadex (Bridion)                      | Restricted to use in the main operating room and main PACU  |
|                                   | Prothrombin complex concentrate (Kcentra) | Restricted to reversal of warfarin, rivaroxaban, or apixaban in patients with life-threatening intracranial bleeding  |
|                                   | Acetylcysteine (Acetadote/Mucomyst)       | <ul style="list-style-type: none"> <li>• Nebulization restricted to use by intensivists (Trauma, CCMS, LSU ICU) in patients who failed nebulized 3% NaCl</li> <li>• Oral and IV acetylcysteine is restricted to treatment of APAP toxicity only. Other indications must be approved by clinical team</li> </ul> |
| Cardiac medications               | Dronedarone (Multaq)                      | Restricted to continuation of home medication   |
|                                   | Ethacrynic acid (Edecerin)                | Restricted to use in patients with heart failure who have hypersensitivity to sulfa-containing drugs  |
| Sclerosing Agent                  | Sodium Tetradecyl Sulfate (Sotradecol)    | Restricted to treatment of severe bleeding secondary to esophageal varices that has failed other treatment modalities. Reserved mainly for pediatric patients at OLOL but can also be used for adult patients   |
| Antiepileptics                    | Clobazam (Onfi)                           | Restricted to use as continuation of home med   |
| H <sub>2</sub> Antagonist and PPI | Ranitidine (Zantac)                       | Restricted to pediatric patients  |
|                                   | Esomeprazole (Prilosec)                   | Restricted to pediatric patients  |
| Malignant hyperthermia treatment  | Dantrolene (Ryanodex)                     | Ryanodex: restricted to malignant hypothermia carts and for first dose treatment for malignant hypothermia; subsequent doses should be administered using IV dantrium   |
| Inhalers                          | Fluticasone inhalation (Flovent)          | Flovent is restricted to pediatric patients   |

|               |   |   |
|---------------|---|---|
|               | Fluticasone-salmeterol (Advair)             | <ul style="list-style-type: none"> <li>• Advair HFA is restricted to kids &lt;12 yrs and unable to use Breo</li> </ul>                            |
| Miscellaneous | Complement C1 esterase inhibitor (Berinert) | <ul style="list-style-type: none"> <li>• Restricted to use in patients with confirmed or suspected hereditary angioedema only</li> </ul>          |
|               | Calcitonin IM (Miacalcin)                   | Restricted to 2 doses in patients with symptomatic hypercalcemia and uncorrected calcium level of at least 13                                     |
|               | Levothyroxine IV (Synthroid)                | Restricted to patients with myxedema coma, LOPA patients, or if a patient is NPO for a prolonged period.  |
|               | Tolvaptan (Samsca)                          | Restricted to treatment of euvolemic and hypervolemic hyponatremia  |
|               | Meningococcal group B vaccine (Bexsero)     | Restricted for patients at least 10 years old with either of the following: initiation of eculizumab therapy or anatomical or functional asplenia |
|               | Corticotropin (Acthar Gel)                  | Restricted to treatment of infantile spasms   |